

# Arid Zone Trees



## *Acacia estrophiolata* Ironwood



### Horticultural Qualities *Acacia estrophiolata* Ironwood

**Foliage:** Evergreen  
**Mature Height:** 25' - 45'  
**Mature Width:** 15' - 25'  
**Growth Rate:** Slow  
**Hardiness:** 15 degrees F  
**Exposure:** Full Sun  
**Leaf Color:** Pale Green  
**Shade:** Dense  
**Flower Color:** Cream to Pale Yellow  
**Flower Shape:** Ball  
**Flower Season:** Spring  
**Thorns:** No  
**Box Sizes Produced:** Not available  
**Propagation Method:** Seed

[www.aridzonetrees.com](http://www.aridzonetrees.com)

Arid Zone Trees, P. O. Box 167, Queen Creek, AZ 85242, Phone 480-987-9094 e-mail: [info@aridzonetrees.com](mailto:info@aridzonetrees.com)

## *Acacia estrophiolata* Ironwood

*Acacia estrophiolata* (Ironwood) is a graceful tree from Australia with a spreading crown and when mature a weeping habit. As a young tree it has rigid branches and short straight leaves (phyllodes) in clusters. The trees characteristics morph as it matures with a weeping pendulous branching structure and the leaves are longer and not clustered. Its pale green foliage blends well with our southwestern native habitat. A heavy bloom in the spring of pale yellow ball flowers brightens the landscape. This is a useful long-lived drought and frost tolerant shade and street tree when planted in well drained soils in arid landscapes. This tree experienced no damage at 15 degree F. (below 25 degree F. for 11 hours) in our nursery.

Additional research needs to be collected.

### Cultural Practices

Foster the development of a more dispersed root system and reduce the risk of wind throw by arranging irrigation emitters at varying distances from the trunk to encourage roots to "seek out" water and nutrients. Irrigation emitter arrangement along with other information on irrigations practices for desert trees can be found at [www.aridzonetrees.com](http://www.aridzonetrees.com) and click on the FAQ link.

Prune as needed to reinforce the structure and form of the tree. Periodic thinning is the most desirable method of pruning. Avoid hedging or heading back desert species, as this will only stimulate excessive branching. Do not remove more than 30% of the canopy during the summer as this can lead to sunburn injuries that can later be invaded by wood boring insects. Always use clean, sharp tools that are cleaned regularly in a 10% solution of bleach. For detail pruning guide see [www.aridzonetrees.com](http://www.aridzonetrees.com) and click on the FAQ interactive button.

Periodically insect pests can be a problem on some desert trees. On young trees, insect infestation can slow typical seasonal growth. Inspect trees during the growing season for common garden sucking insects such as aphids, thrip, whiteflies or psyllids. During dry months, (May and June) in dusty conditions, spider mites can appear. Monitor for infestation and apply controls as needed. Spray applications of water or water and Safer Soap give short-term control (3 to 7 days) for small insect population. For heavy infestation or longer control use federally registered insecticides. A contact insecticide application will kill existing adults. An application with a systemic soil drench will provide 8 to 12 weeks control for any post application insect hatchings or migration of insects. Before using pesticide for the first time or on new plants or cultivar, treat a few plants and check for phytotoxicity. **Always read label and follow label instruction before using pesticides. For pesticide control recommendations contact a licensed pest control advisor.**

[www.aridzonetrees.com](http://www.aridzonetrees.com)